

Research Application Summary

**Effect of banana peels on chicken weight gain and egg production in the urban and peri-urban areas of Aksum City, Ethiopia**

Haileselassie, G.A.<sup>1</sup>, Gebrechristos, S.G.<sup>2</sup> & Wasonga, O.V.<sup>3</sup>

<sup>1</sup>Department Dryland Agriculture and Natural Resources Management, IGAD Sheikh Technical Veterinary School, Terra Nouva East Africa, P. O. Box 74916 - 00200, Nairobi, Kenya

<sup>2</sup>Department of Animal Science and Technology, Aksum University, P. O. Box 1010, Tigray, Ethiopia

<sup>3</sup>Department of Land Resource Management and Agricultural Technology, University of Nairobi, P. O. Box 29053, Nairobi, Kenya

**Corresponding author:** [hailish746990@gmail.com](mailto:hailish746990@gmail.com)

---

**Abstract**

High feed cost of running urban-animal-agriculture and huge city wastes are challenges of the growing towns in modern Ethiopia. Egg production microenterprise is steadily growing due to its low initial investment and ready market in urban and peri-urban areas. However, cost of egg production is high due to the rising cost of maize, the main feed ingredient in poultry rations. Using banana peels, which forms a great proportion of city waste in Ethiopia, is seen as a way of not only reducing the waste, but also as an alternative low-cost feed supplement that could help enhance egg production by small-holder farmers. This study looked at the effect of feeding banana-peels-powder on weight-gain, egg-production and profit from rearing improved poultry breeds. Banana peels were air-dried, ground and used to re-formulate existing feed rations. The control was the poultry ration formulated with 52% maize and 48% other ingredients. Then four rations were formulated with 13%, 26%, 39% and 52% banana peels. One hundred, 5-months old Bovans Brown chicken breed were clustered into four blocks based on weight categories. Five treatments were applied randomly to the blocks and replicated five times in a RCBD cage battery system. The chickens were fed with 120g/chicken/day while water was provided *ad libitum*. Chicken weight and eggs produced were recorded every two weeks and daily, respectively. Whereas weight gain and egg production declined with increase in proportion of banana peels, 25% replacement resulted in a comparable daily weight gain (26.42 gm) to the standard chicken ration (26.76 gm). Increase of banana peels resulted in reduction of weight gain at 17.78, 16.67 and 15.11 gm/ day and egg production at 0.58, 0.4 and 0.22/day/chicken for 50, 75 and 100% replacements respectively. We conclude that 25% banana peels powder in maize-based feed ration gives optimum weight gain and egg production in chickens.

Key words: Banana peels formulation, Bovans brown, feed ratio formulation, replacement, waste

**Résumé**

Le coût élevé des aliments pour l'agriculture urbaine et les immenses déchets urbains sont des défis pour les villes en pleine croissance de l'Éthiopie moderne. La micro-entreprise de production d'œufs est en croissance constante en raison de son faible investissement initial et de son marché prêt dans les

zones urbaines et périurbaines. Cependant, le coût de la production d'œufs est élevé en raison de la hausse du coût du maïs, principal ingrédient alimentaire des rations de volaille. Utilisation de pelures de banane, qui constitue une grande partie des déchets urbains en Éthiopie, est considérée comme un moyen non seulement de réduire les déchets, mais aussi comme un complément alimentaire alternatif à faible coût qui pourrait aider à améliorer la production d'œufs par les petits agriculteurs. Cette étude a examiné l'effet de l'alimentation en poudre de peau de banane sur le gain de poids, la production d'œufs et les bénéfices de l'élevage de races de volaille améliorées. Les pelures de banane ont été séchées à l'air, broyées et utilisées pour reformuler les rations alimentaires déclarées. Le témoin était la ration de volaille formulée avec 52% de maïs et 48% d'autres ingrédients. Ensuite, quatre rations ont été formulées avec 13%, 26%, 39% et 52% de pelures de banane. Cent races de poulets Bovans Brown âgées de 5 mois ont été regroupées en quatre blocs en fonction des catégories de poids. Cinq traitements ont été appliqués au hasard aux blocs et reproduits cinq fois dans un système de batterie à cage RCBD. Les poulets ont été nourris avec 120 g / poulet / jour tandis que de l'eau était fournie ad libitum. Le poids des poulets et les œufs produits ont été enregistrés toutes les deux semaines et publiées, respectivement. Alors que le gain de poids et la production d'œufs ont diminué avec l'augmentation de la proportion de pelures de banane, un remplacement de 25% a requis un gain de poids quotidien comparable (26,42 g) à la ration de poulet standard (26,76 g). L'augmentation des pelures de banane a suivi une réduction du gain de poids à 17,78, 16,67 et 15,11 g / jour et de la production d'œufs à 0,58, 0,4 et 0,22 / jour / poulet pour 50, 75 et 100% de remplacements respectivement. Dans une ration alimentaire à base de maïs donne un gain de poids et une production d'œufs optimaux chez les poulets.

Mots clés : formulation de pelures de banane, Bovans brun, formulation de rapport d'alimentation, remplacement, déchets

---

## Introduction

Banana has many common names (Espino *et al.*, 1991) and belongs to the genus *Musa* spp. and is a popular fruit worldwide. Banana peels are the outer envelopes and by-product of household consumption and banana processing which is widely available in populated areas, especially in the vicinity of processing plants. Banana peels contain 6-9% crude protein and 20-30% NDF on dry matter basis (Emaga *et al.*, 2011). Green plantain peels contain 40% starch that is fully transformed into sugars after ripening, while banana peels contain much less starch (about 15%) when green and end up with 30% free sugars when ripe. Lignin content also increases with ripening (from 7 to 15% DM). Plantain peels contain less fibre than bananas (Emaga *et al.*, 2011).

Banana peels are widely used by smallholders as complementary feeds for cattle and small ruminants in the tropics (Onwuka *et al.*, 1997; Emaga *et al.*, 2011). In a study done by FAS *et al.* (2015) in Nigeria, feeding broilers on diets containing 10% banana peels caused no adverse effect on performance. Moreover, Aregheore (1998) in his research in Nigeria reported that dry ripe plantain peels can replace up to 100% maize without adverse effects on growth and performance of goats, and were found an economical source of carbohydrates. Nambi-Kasozi *et al.* (2014) also reported that dairy cows fed with 40-60% banana peels in their daily ration in Uganda showed higher dry matter intake than 0-20% inclusion with no significant difference on daily weight gain and milk yield. According to Oyedeji (2015), highest weight gain per bird (1432 g) and the best feed/gain ratio (2.58) were recorded among broilers fed 50% ripe banana peels in an experiment conducted in Benin University, Nigeria. In addition, live weight and feed conversion efficiency have been found to significantly increase

when chicken were fed on 10% banana peel meal. However, growth performance decreased when the inclusion rate exceeded 10% (Sabutan, 1996).

Banana peels form a large proportion of household and therefore Municipal wastes in towns in the Tigray region of Ethiopia. For instance, according to the Trade, Industry and Urban Development Office (Personal communication) out of the approximately 10 tons of garbage generated in Aksum Town daily, a significant amount is banana peels and more than 30% of garbage rots uncollected on pavements, streets, sewerage outlets and water channels due to the high costs of collection and disposal. On the other hand, scarcity and high cost of animal feed lowers profit margin of small and microenterprises in the region. However, despite the availability of banana peels, empirical evidence is scarce on its potential as complimentary animal feed that would provide low-cost alternative for the small holder farmers in Tigray, especially those living in the urban and peri-urban areas. This study, therefore, aimed at determining the effect of different proportions of banana peels on feed intake, weight gain, egg production, as well as cost saved in a poultry enterprise.

### Materials and methods

The study was conducted in Aksum University located in the Central zone of Tigray, 784 km North to Addis Ababa, Ethiopia's capital city. Banana peels were collected from Aksum town. Rations were formulated using different proportions of sundried banana peels powder mixed with a pre-formulated feed. Banana peels were sundried, ground and mixed with wheat bran (8%), middling (8%), meat and bone meal (8%), molasses (2%), protein sources (15%), minerals (8%) to form rations containing 0%, 13%, 26%, 39% and 52% banana peel powder (Table 1).

**Experimental design.** A total of 100 Bovance brown chickens of five months of age were used for the experiment. The birds were classified into four blocks based on the weight category. The four classes were randomly allocated to the five treatments and replicated five times in a RCBD cage battery system.

**Data collection.** A feeding trial was conducted for a period of 45 days with an acclimatization period of 15 days before starting data collection. The experimental birds were weighed before and then after every two weeks for 6 weeks, while egg production was recorded daily. During the experiment, 120g of feed/day/chicken was provided at 9:00 a.m. in the morning and 5:00 p.m. in the evening, while water was provided ad libitum. Fresh water made available with individual waterers ad lib throughout the trial period. All necessary health care services were provided to the experimental chicken.

**Comparative cost saving analysis.** Cost savings were analyzed using the method by FAS *et al.* (2015). Feed cost was obtained through summing the cost of ingredients of the total feed in diet composition. Whereas, the cost of the feed per weight gain (kg) was calculated by multiplying the cost with feed conversion ratio. The cost of feed per egg was calculated through dividing the total number of experimental days (45) by the total number of eggs obtained within the 45 experimental days multiplied by cost of daily feed intake. The feed cost savings was calculated by subtracting supplementation by banana peels (Final costs calculated from cost incurred to collect banana peels) from costs without supplementation by banana peels (initial costs) multiplied by 100 as follows:

$$\text{Feed cost savings} = (\text{Initial cost} - \text{Final cost}) \times 100$$

The initial cost was also computed by multiplying cost of feed per kg by the amount (g) of daily feed

intake as follows:

Cost of daily feed intake (per bird) = Cost of feed/kg x Quantity of daily feed intake

**Table 1. Ingredients and proportions (%) of the experimental diets**

No.	Ingredient	Proportions of each ingredient (%) along the five diets				
		Diet 1 (Control)	Diet 2	Diet 3	Diet 4	Diet 5
1	Noug cake	6.0	6.0	6.0	6.0	6.0
2	Corn	52.0	39.0	26.0	13.0	0.0
3	Ground banana peels Ratio	0.0	13.0	26.0	39.0	52.0
4	Wheat bran	8.0	8.0	8.0	8.0	8.0
5	Wheat Middling	8.0	8.0	8.0	8.0	8.0
6	Limestone	8.0	8.0	8.0	8.0	8.0
7	Fishmeal	2.0	2.0	2.0	2.0	2.0
8	Molasses	2.0	2.0	2.0	2.0	2.0
9	Salt	0.2	0.2	0.2	0.2	0.2
10	Lysine	0.08	0.08	0.08	0.08	0.08
11	Methionine	0.028	0.028	0.028	0.028	0.028
12	Meat and bone meal	8.0	8.0	8.0	8.0	8.0
13	Layer premix	0.6	0.6	0.6	0.6	0.6
14	Soya bean	5.0	5.0	5.0	5.0	5.0
15	Colin Chloride	0.025	0.025	0.025	0.025	0.025

**Data analysis.** SAS (2008) JMP release 8 Statistical package was used to analyze the data. Analysis of variance was used to determine if there were significant differences in weight gain and egg production between the treatments. The means in weight gain and egg production were separated using LSD at  $P \leq 0.05$ .

## Results and discussion

**Weight gain.** The average weight gain and egg production of the layers decreased as the proportion of banana peels in the chickens' diet increased (Table 2). The mean weight gain of birds fed on the ration without banana peels powder (1204 gm) was the highest, whereas, the feed ration with 100% banana peels (680 gm) had the lowest weight gain.

**Table 2. Mean weight gain and egg production of layers by treatment**

Variable	Treatment					SE
	T1	T2	T3	T4	T5	
Weight gain (g)	1204b	1189b	800a	750a	680a	±50.6
Egg production (No)	36c	32c	26b	18a	10a	±2.2

<sup>a</sup> Rows treatments means with different superscript are significantly different ( $P < 0.05$ )

T1-T5: see Table 1

Replacement of maize by banana peels up to a proportion of 25% did not affect the net weight gain significantly. Previous study conducted by Widjastuti and Hernawan (2012) in Indonesia indicated that replacement of maize with 20% of banana peels does not affect the net weight of broilers.

**Egg production.** Replacement of maize by banana peels exceeding 75% significantly reduced egg production (Table 2), whereas replacing 25% of maize by banana peels did not cause any significant change in egg production. In addition, replacing maize by 50% banana peels resulted in slight decline in egg production.

**Comparative cost analysis of feed ingredients.** The analysis of feed cost per kg indicated that T1 was most expensive (Birr 10.6) of the four rations (Table 3). The cost of formulating T2, T3, T4 and T5 were Birr 9.98, 9.74, 8.87 and 7.78 per kg ration, respectively. These results show that the cost of chicken feed decreased with the increase of proportion of banana peels in the ration. The decrease in feed cost was associated with replacement of cheap source of feed staff (banana peels) instead of the alternative and comparatively expensive feed ingredient (maize) in the balanced poultry ration. Similar results have been reported from experiments conducted on broilers by FAS *et al.* (2015) in Indonesia. The cost of feed intake per chicken was significantly higher ( $p < 0.05$ ) for T1 (Birr 1.45) than all other treatments.

The results show that T5 had the highest (Birr 34.12) cost of feed per body weight gain, while the lowest cost was incurred for T1 (Birr 29.45) and T2 (Birr 29.33). The high cost incurred per body weight gain for T4 and T5 might be associated with low feed conversion ratio as proportion of banana peels in the chickens diet increased. This result is in agreement with the findings of FAS *et al.* (2015) who reported that feed efficiency ratio decreased as the level of peel meal inclusion increased from 0%, 10%, 20% and 30% in broilers diet in Nigeria. Similarly, the analysis of feed cost per egg production revealed that the highest cost incurred per single egg produced was observed in T5 (Birr 4.45). The expense incurred per egg for T4, T3, T2, were Birr 2.825, Birr 2.05, Birr 1.77 and Birr 1.87, respectively. The low cost for T2 could be linked to the low cost incurred for the feed without any significant decrement in egg laying observed for feed ration with 25% replacement of corn by banana peels. Similarly, the low cost of feed/kg weight gain for T2 may be associated with low feed cost incurred as a result of 25% replacement of banana peels with maize in the balanced ration without any significant impact in body weight gain as compared to control group. The analysis made on feed cost saving revealed that T5 had the highest (Birr 26.6) feed cost saving compared to Birr 16.32, Birr 8.11, Birr 5.84 for T4, T3 and T2, respectively. This is attributed to the low cost of banana peels with the only requirement of labor cost for collection, drying, grinding and packaging as compared to maize in the balanced feed of poultry ration formulation.

**Table 3. Cost savings benefit analysis of banana peel replaced poultry diet (Birr)**

Parameter	T1	T2	T3	T4	T5	SD
Cost of feed/kg of diet (Birr)	10.6a	9.98b	9.74ab	8.87 ab	7.78c	±0.689
Cost of feed intake per chicken (Birr)	1.45 a	1.26a	1.19b	1.13b	0.99c	±0.443
Cost of feed/kg weight gain (Birr)	29.45a	29.33a	30.12b	32.12ab	34.12c	±1.208
Cost of feed/egg (Birr)	1.812a	1.77a	2.05b	2.825ab	4.45c	±1.117
Feed cost savings (%)	-	5.84c	8.11c	16.32b	26.6a	±0.089

Means within same row with different superscripts differ significantly ( $P < 0.05$ )

## Conclusion

The results show that whereas weight gain and egg production declined with increase in proportion of banana peels powder in diet of chickens, the feed ration with 13% banana peel (25% maize replacement) did not adversely affect weight gain, egg production, incurred lower cost per kg of weight gain and per egg produced. The cost savings was found to be highest for the 100% replacement of maize with banana peels due to low cost of feed. On the other hand replacement of maize by over 75% banana peels adversely affected egg laying ability of birds and showed high cost per weight gain and per egg produced. We conclude that banana peels powder can replace up to 25% of corn in chicken diet without causing adverse effect on their performance. Inclusion beyond 13% of the whole ration of chicken is not recommended.

## Acknowledgements

We thank Tigrai Science and Technology Agency for the financial grant to conduct this study. We also acknowledge the Director of Sefho Business Enterprise of Aksum University and Aksum University Research Directorate for their support during the study period. This paper is a contribution to the Sixth Africa Higher Education Week and RUFORUM 2018 Biennial Conference.

## References

- Abel, F.A.S., Adeyemi, O.A., Oluwole, O.B., Oladunmoye, O.O., Ayo-Ajasa, O.Y. and Anuoluwatele, J.O. 2015. Effects of treated banana peel meal on the feed efficiency, digestibility and cost effectiveness of broiler chickens diet. *J. Vet. Sci. Anim. Husb* 3 (1): 101-107.
- Aregheore, E.M. 1998. A note on the nutritive value of dry ripe plantain peels as a replacement of maize for goats. *J. Anim. Feed Sci* 7 (1): 55-62.
- Emaga, T.H., Bindelle, J., Agneensens, R., Buldgen, A., Wathelet, B. and Paquot, M. 2011. Ripening influences banana and plantain peels composition and energy content. *Tropical Animal Health and Production* 43 (1): 171-177.
- Nambi-Kasozi, J., Sabiiti, E.N., Bareeba, F.B. and Sporndly, E. 2014. Effect of feeding varying levels of banana peelings supplemented with maize bran, cotton seed cake and *Gliricidia sepium* on the performance of lactating dairy cows. *African Journal of Agricultural Research* 9 (8): 722-27.
- Onwuka, C.F.I., Adetiloye, P.O. and Afolami, C.A. 1997. Use of household wastes and crop residues in small ruminant feeding in Nigeria. *Small Ruminant Research* 24 (3): 233-37. <http://linkinghub.elsevier.com/retrieve/pii/S0921448896009534>.
- Oyedeji, J.O., Olomu, J.M., Godspower, S.A. and Ojero, O.V. 2015. Effects of graded levelsof ripe and unripe plantain peel meals onperformance, organ weights and bloodparameters of broiler chickens. *NJAFE* 11: 28-33.
- Sabutan, M.G.I. 1996. Banana peelings help broilers grow. *Misset World Poultry* 12 (6): 59-61.
- SAS. 2008.JMP Release 8. Cary, NC, SAS Institute Inc.
- Widjastuti, T. and Hernawan, E., 2012. Utilizing of banana peel (*Musa sapientum*) in the ration and its influence on final body weight, percentage of carcass and abdominal fat on broilers under heat stress condition. *Zootehnie* 57: 104-109.